

# Practical work 5

## Student-Test

This example teaches you how to perform a t-Test in Excel. The t-Test is used to test the null hypothesis that the means of two populations are equal.

Below you can find the study hours of 6 female students and 5 male students.


$$H_0: \mu_1 - \mu_2 = 0$$

$$H_1: \mu_1 - \mu_2 \neq 0$$


	A	B	C
1	Female	Male	
2	26	23	
3	25	30	
4	43	18	
5	34	25	
6	18	28	
7	52		
8			

To perform a t-Test, execute the following steps.


1. First, perform an **F-Test** to determine if the variances of the two populations are equal. This is not the case.
2. On the Data tab, in the Analysis group, click Data Analysis.



3. Select t-Test: Two-Sample Assuming Unequal Variances and click OK.



4. Click in the Variable 1 Range box and select the range A2:A7.
5. Click in the Variable 2 Range box and select the range B2:B6.
6. Click in the Hypothesized Mean Difference box and type 0 ( $H_0: \mu_1 - \mu_2 = 0$ ).
7. Click in the Output Range box and select cell E1.



8. Click OK.

Result:

E	F	G
t-Test: Two-Sample Assuming Unequal Variances		
	<i>Variable 1</i>	<i>Variable 2</i>
Mean	33	24.8
Variance	160	21.7
Observations	6	5
Hypothesized Mean Difference	0	
df	7	
t Stat	1.47260514	
P(T<=t) one-tail	0.092170202	
t Critical one-tail	1.894578605	
P(T<=t) two-tail	0.184340405	
t Critical two-tail	2.364624252	

Conclusion: We do a two-tail test (inequality). If  $t \text{ Stat} < -t \text{ Critical two-tail}$  or  $t \text{ Stat} > t \text{ Critical two-tail}$ , we reject the null hypothesis. This is not the case,  $-2.365 < 1.473 < 2.365$ . Therefore, we do not reject the null hypothesis. The observed difference between the sample means (33 - 24.8) is not convincing enough to say that the average number of study hours between female and male students differ significantly.

Ex: Apply the above in series 6.